# antibodies.com

## Datasheet for ABIN7321079 DDR1 Protein (Fc Tag)

Image



#### Overview

Quantity:	100 µg
Target:	DDR1
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This DDR1 protein is labelled with Fc Tag.

#### Product Details

Purpose:	Recombinant Rat DDR1 Kinase/CD167 Protein (Fc Tag)
Sequence:	Met1-Thr413
Characteristics:	A DNA sequence encoding the rat DDR1 (Q6MG19) (Met1-Thr413) was expressed with the Fc region of human IgG1 at the C-terminus.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per $\mu$ g of the protein as determined by the LAL method

#### Target Details

Target:	DDR1
Alternative Name:	DDR1 (DDR1 Products)
Background:	Background: Discoidin domain receptor family, member 1 (DDR1), also known as or CD167a (cluster of differentiation 167a), and Mammary carcinoma kinase 10 (MCK10), belongs to a
	subfamily of tyrosine kinase receptors with an extracellular domain homologous to


Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7321079 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

	Dictyostellium discoideum protein discoidin 1. Receptor tyrosine kinases play a key role in the
	communication of cells with their microenvironment. These kinases are involved in the
	regulation of cell growth, differentiation and metabolism. Expression of DDR1/MCK10/CD167 is
	restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In
	addition, it has been shown to be significantly overexpressed in several human tumors.
	DDR1/MCK10/CD167 plays an important role in regulating attachment to collagen, chemotaxis,
	proliferation, and MMP production in smooth muscle cells. DDR1 functions in a feedforward
	loop to increase p53 levels and at least some of its effectors. Inhibition of DDR1 function
	resulted in strikingly increased apoptosis of wild-type p53-containing cells in response to
	genotoxic stress through a caspase-dependent pathway.
	Synonym: DDR1,MCK10
Molecular Weight:	70.9 kDa
UniProt:	Q6MG19
Pathways:	RTK Signaling, Smooth Muscle Cell Migration

### Application Details

Comment:	71 kDa
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN7321079 | 09/09/2023 | Copyright antibodies-online. All rights reserved.



Western Blotting

Image 1.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7321079 | 09/09/2023 | Copyright antibodies-online. All rights reserved.