

Datasheet for ABIN7321101

**JAM2 Protein (Fc Tag)**[Go to Product page](#)**1** Image

## Overview

Quantity:	100 µg
Target:	JAM2
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This JAM2 protein is labelled with Fc Tag.

## Product Details

Purpose:	Recombinant Rat JAM-B/CD322 Protein (Fc Tag)
Sequence:	Met1-Asn236
Characteristics:	A DNA sequence encoding the rat JAM2 (Q3MHC0) (Met1-Asn236) was expressed, fused with the Fc region of human IgG1 at the C-terminus.
Purity:	> 90 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method

## Target Details

Target:	JAM2
Alternative Name:	JAM-B/CD322 ( <a href="#">JAM2 Products</a> )
Background:	Background: Junctional adhesion molecule B (JAM-B), also known as Junctional adhesion molecule 2 (JAM2), Vascular endothelial junction-associated molecule (VE-JAM), and CD322, is a single-pass type I membrane protein which belongs to the immunoglobulin superfamily. It is

## Target Details

prominently expressed on high endothelial venules. expression to be restricted to the high endothelial venule of tonsil and lymph nodes. The localization to the endothelium of arterioles in and around inflammatory and tumor foci. JAM-B can function as an adhesive ligand for the T cell line J45 and can interact with GM-CSF/IL-4-derived peripheral blood dendritic cells, circulating CD56(+) NK cells, circulating CD56(+)CD3(+) NK/T cells, and circulating CD56(+)CD3(+)CD8(+) cytolytic T cells. JAM-2 is expressed on high endothelial venules (HEVs) in human tonsil and on a subset of human leukocytes, suggesting that JAM-2 plays a central role in the regulation of transendothelial migration. It binds to very late activation antigen (VLA)-4, a leucocyte integrin that contributes to rolling and firm adhesion of lymphocytes to endothelial cells through binding to vascular cell adhesion molecule (VCAM)-1. JAM-B appears to contribute to leucocyte extravasation by facilitating not only transmigration but also rolling and adhesion. JAM-B acts as an adhesive ligand for interacting with a variety of immune cell types and may play a role in lymphocyte homing to secondary lymphoid organs.

Synonym: JAM2,JAM-2

Molecular Weight:	50.3 kDa
-------------------	----------

UniProt:	<a href="#">Q3MHCO</a>
----------	------------------------

## Application Details

Restrictions:	For Research Use only
---------------	-----------------------

## Handling


Format:	Lyophilized
---------	-------------

Reconstitution:	Please refer to the printed manual for detailed information.
-----------------	--

Buffer:	Lyophilized from sterile PBS, pH 7.4
---------	--------------------------------------

Storage:	4 °C,-20 °C,-80 °C
----------	--------------------

Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
------------------	---



Western Blotting

Image 1.