

### Datasheet for ABIN7321108

# **PROCR Protein (His tag)**





#### Overview

Quantity:	100 μg
Target:	PROCR
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This PROCR protein is labelled with His tag.

### **Product Details**

Purpose:	Recombinant Rat EPCR Protein (His Tag)
Sequence:	Met1-Ser213
Characteristics:	A DNA sequence encoding the rat PROCR (Met1-Ser213) was expressed with a polyhistidine tag at the C-terminus.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin Level:	$<$ 1.0 EU per $\mu g$ of the protein as determined by the LAL method

## Target Details

Target:	PROCR
Alternative Name:	EPCR (PROCR Products)
Background:	Background: Endothelial protein C receptor (EPCR), also known as activated protein C receptor (APC receptor) or PROCR, is a receptor for Protein C. Protein C plays an important role in many
	metabolism processes in humans and other animals after activated by binding to Endothelial

protein C receptor (EPCR). Because of the EPCR is found primarily on endothelial cells (cells on the inside of blood vessels), activated protein C is found maily near endothelial cells. Protein C is pleiotropic, with two main functions: anticoagulation and cytoprotection. Which function will be performed depend on whether or not protein C remains bind to EPCR after activated. The anticoagulation occurs when it does not. In this case, protein C functions as an anticoagulant by irreversibly proteolytically inactivating Factor Va and Factor VIIIa, turning them into Factor Vi and Factor VIIIi respectively. When still bound to EPCR, activated protein C performs its cytoprotective effects, acting on the effector substrate PAR-1, protease-activated receptor-1. To a degree, APC's anticoagulant properties are independent of its cytoprotective ones, in that expression of one pathway is not affected by the existence of the other.

Synonym: PROCR

Molecular Weight:

23.2 kDa

UniProt:

Q4V8I1


#### **Application Details**

Restrictions:

For Research Use only

#### Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.



## **Western Blotting**

Image 1.