



[Go to Product page](#)

Datasheet for ABIN7321137
CNTN3 Protein (Fc Tag)

1 Image

Overview

| | |
|-------------------------------|---|
| Quantity: | 100 µg |
| Target: | CNTN3 |
| Origin: | Rat |
| Source: | HEK-293 Cells |
| Protein Type: | Recombinant |
| Purification tag / Conjugate: | This CNTN3 protein is labelled with Fc Tag. |

Product Details

| | |
|------------------|--|
| Purpose: | Recombinant Rat Contactin 3/CNTN3 Protein (Fc Tag) |
| Sequence: | Met1-Gly1001 |
| Characteristics: | A DNA sequence encoding the rat CNTN3 (Met1-Gly1001) was expressed with the Fc region of human IgG1 at the C-terminus. |
| Purity: | > 85 % as determined by SDS-PAGE |
| Endotoxin Level: | < 1.0 EU per µg of the protein as determined by the LAL method |

Target Details

| | |
|-------------------|--|
| Target: | CNTN3 |
| Alternative Name: | Contactin 3/CNTN3 (CNTN3 Products) |
| Background: | Background: Contactins are a subgroup of molecules belonging to the immunoglobulin superfamily that are expressed exclusively in the nervous system. The subgroup consists of six members: Contactin-1, Contactin-2(TAG-1), Contactin-3(BIG-1), BIG-2, Contactin-5(NB-2) and |

Target Details

NB-3. Since their identification in the late 1980s, Contactin-1 and Contactin-2 have been studied extensively. Axonal expression and the neurite extension activity of Contactin-1 and Contactin-2 attracted researchers to study the function of these molecules in axon guidance during development. Contactin-1 and Contactin-2 have come to be known as the principal molecules in the function and maintenance of myelinated neurons. In contrast, the function of the other four members of this subgroup remained unknown until recently. Contactin-3, also known as CNTN3 (BIG-1 in rat and PANG in mouse), is a GPI-linked glycoprotein that is expressed on cerebellar Purkinje cells, amygdaloid and thalamic neurons and olfactory granule cells. In the brain, Contactin-3 is expressed in frontal lobe, occipital lobe, cerebellum and amygdala. Contactin-3 contains 4 fibronectin type-III domains and 6 Ig-like C2-type (immunoglobulin-like) domains. Human Contactin-3 shares 92 % aa identity with mouse Contactin-3. The exact function of Contactin-3 is unclear. Contactin-3 may mediate cell-cell interaction and may promote neurite outgrowth.

Synonym: CNTN3

Molecular Weight: 134.8 kDa

NCBI Accession: [NP_062202](#)

Application Details

Restrictions: For Research Use only

Handling


Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile PBS, pH 7.4

Storage: 4 °C,-20 °C,-80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.