antibodies -online.com





OLR1 Protein (Fc Tag)





Go to Product page

Overview

Quantity:	100 μg
Target:	OLR1
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This OLR1 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Rat OLR1/LOX1 Protein (Fc Tag)
Sequence:	Leu60-Gln364
Characteristics:	A DNA sequence encoding the rat OLR1 (070156) (Leu60-Gln364) was expressed with Fc region of human IgG1 at the N-terminus.
Purity:	(90.4+8.5) % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method

Target Details

Target:	OLR1
Alternative Name:	OLR1/LOX1 (OLR1 Products)
Background:	Background: Oxidized low-density lipoprotein receptor 1 (Ox-LDL receptor 1 or OLR1), also
	known as lectin-type oxidized LDL receptor 1 (LOX1), is a receptor protein that belongs to the C-
	type lectin superfamily. LOX1 is a multi-ligand receptor originally identified as the endothelial

oxidized LDL receptor. OLR1 / LOX1 was isolated from an aortic endothelial cell, and recently it has been discovered in macrophages and vascular smooth muscle cells in artery vessels. The expression of LOX1 is inducted by inflammatory stimuli and oxidative stimuli. This protein binds, internalizes and degrades oxidized low-density lipoprotein. LOX1 may play an important role in the progression of vulnerable carotid plaque and might regulate vulnerable plaque formation in cooperation with MMPs and TIMP-2. In clinical, LOX1 is thought to be involved in the development of atherosclerotic lesions.

Synonym: OLR1;Lox1;Oldlr1

Molecular Weight:

63.9 kDa

UniProt:

070156


Application Details

Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.