antibodies - online.com







REG3A Protein (Fc Tag)





Overview

Quantity:	100 μg
Target:	REG3A
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This REG3A protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Rat REG3A/HIP Protein (Fc Tag)
Sequence:	Met1-Gln174
Characteristics:	A DNA sequence encoding the rat REG3A (BAA04904.1) (Met1-Gln174) was expressed, fused with the Fc region of human IgG1 at the C-terminus.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin Level:	$<$ 1.0 EU per μg of the protein as determined by the LAL method

Target Details

Target:	REG3A
Alternative Name:	REG3A/HIP (REG3A Products)
Background:	Background: Regenerating islet-derived protein 3-alpha, also known as Regenerating islet- derived protein III-alpha, REG-3-alpha, REG3A, and HIP, is secreted protein which contains one
	C-type lectin domain. REG3A is constitutively expressed in intestine, and is a pancreatic

Target Details

secretory protein that may be involved in cell proliferation or differentiation. It is overexpressed during the acute phase of pancreatitis and in some patients with chronic pancreatitis. REG3A and REG1A proteins are both involved in liver and pancreatic regeneration and proliferation. REG3A is also a stress protein involved in the control of bacterial proliferation. REG3A is down-regulated in most primary human gastric cancer cells, and might be useful in the diagnosis of gastric cancer. Additionally, REG3A is a target of beta-catenin signaling in Huh7 hepatoma cells. The REG1A and REG3A are downstream targets of the Wnt pathway during liver tumorigenesis. Synonym: REG3A,Pap2,Reg3

Molecular Weight:

43.4 kDa

Application Details


Restrictions:

For Research Use only

Handling

Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.

Images



Western Blotting

Image 1.