



[Go to Product page](#)

Datasheet for ABIN7321168

## Mbl1 Protein (Fc Tag)

### 1 Image

#### Overview

Quantity:	100 µg
Target:	Mbl1
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Mbl1 protein is labelled with Fc Tag.

#### Product Details

Purpose:	Recombinant Rat MBL1 Protein (Fc Tag)
Sequence:	Met1-Ala238
Characteristics:	A DNA sequence encoding the rat MBL1 (P19999) (Met1-Ala238) was expressed with the Fc region of human IgG1 at the C-terminus.
Purity:	> 85 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method

#### Target Details

Target:	Mbl1
Alternative Name:	MBL1 ( <a href="#">Mbl1 Products</a> )
Background:	Background: Mannose-binding lectin (MBL), also named mannose or mannan-binding protein (MBP), is a C-type lectin which participates in the innate immune system as an activator of the complement system and as opsonin after binding to certain carbohydrate structures on

## Target Details

---

microorganisms and pathogens. Its function appears to be pattern recognition in the first line of defense in the pre-immune host. MBL recognizes carbohydrate patterns found on the surface of a large number of pathogenic micro-organisms including bacteria, viruses, protozoa and fungi. Binding of MBL to a micro-organism results in activation of the lectin pathway of the complement system. Two forms of MBL, MBL-A and MBL-C, were characterized in rodents, rabbits, bovine and rhesus monkeys, whereas only one form was identified in humans, chimpanzees and chickens. The two forms are encoded by two distinct genes named MBL1 and MBL2, which have been identified in many species including the pig. The MBL1 and MBL2 genes encode mannan-binding lectins (MBL) A and C, respectively, that are collagenous lectins (collectins) produced mainly by the liver. The MBL1 gene encodes MBL-A, which has bacteria-binding properties in pigs and rodents but is mutated to a pseudogene in humans and chimpanzees. Deficiency of MBL is probably the most common human immunodeficiency and is associated with an increased risk of mucosally acquired infections including meningococcal disease. MBL could modify disease susceptibility by modulating macrophage interactions with mucosal organisms at the site of initial acquisition.

Synonym: MBL1

---

Molecular Weight: 50.5 kDa

---

UniProt: [P19999](#)

## Application Details

---

Restrictions: For Research Use only

## Handling

---


Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile PBS, pH 7.4

Storage: 4 °C, -20 °C, -80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.



**Western Blotting**

**Image 1.**