# antibodies .- online.com





# F11R Protein (Fc Tag)





$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μg
Target:	F11R
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This F11R protein is labelled with Fc Tag.
Product Details	

Purpose:	Recombinant Rat JAM-A/F11R Protein (Fc Tag)	
Sequence:	Met1-Gly238	
Characteristics:	A DNA sequence encoding the rat F11R () was expressed, fused with the Fc region of human IgG1 at the C-terminus.	
Purity:	> 96 % as determined by SDS-PAGE	
Endotoxin Level:	$<$ 1.0 EU per $\mu g$ of the protein as determined by the LAL method	

#### **Target Details**

Target:	F11R
Alternative Name:	JAM-A/F11R (F11R Products)
Background:	Background: Junctional adhesion molecule-A (JAM-A), also known as F11 receptor (F11R) or Cluster of Differentiation 321 (CD321), is a transmembrane protein expressed at tight junctions
	of epithelial and endothelial cells, as well as on circulating leukocytes. JAM-A protein serves as

#### **Target Details**

a serotype-independent receptor for mammalian orthoreoviruses (reoviruses). It is also a ligand for the integrin LFA1, involves in leukocyte transmigration. As a cell adhesion molecule of the immunoglobulin superfamily, JAM-A protein involves in platelet adhesion, secretion and aggregation, and plays a crucial role in inflammatory thrombosis and atherosclerosis. In addition, it may be a potential therapeutic target for breast cancer.

Synonym: F11R,Jam1

Molecular Weight: 49.9 kDa

UniProt: Q9JHY1

Pathways: Cell-Cell Junction Organization


### **Application Details**

Restrictions: For Research Use only

#### Handling

Format:	Lyophilized	
Reconstitution:	Please refer to the printed manual for detailed information.	
Buffer:	Lyophilized from sterile PBS, pH 7.4	
Storage:	4 °C,-20 °C,-80 °C	
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.	

Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.



## **Western Blotting**

Image 1.