

Datasheet for ABIN7321195

Ectodysplasin A2 Receptor Protein (EDA2R) (Fc Tag)[Go to Product page](#)**1** Image

Overview

Quantity:	100 µg
Target:	Ectodysplasin A2 Receptor (EDA2R)
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This Ectodysplasin A2 Receptor protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Rat XEDAR/EDA2R Protein (Fc Tag)
Sequence:	Met2-Glu137
Characteristics:	A DNA sequence encoding the rat EDA2R (D3ZAX4) (Met2-Glu137) was expressed, fused with the Fc region of human IgG1 at the C-terminus.
Purity:	> 94 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method

Target Details

Target:	Ectodysplasin A2 Receptor (EDA2R)
Alternative Name:	XEDAR/EDA2R (EDA2R Products)
Background:	Background: Tumor necrosis factor receptor superfamily member 27, also known as X-linked ectodysplasin-A2 receptor, EDA-A2 receptor, EDA2R, XEDAR and TNFRSF27, is a single-pass type I II membrane protein. TNFRSF27 / EDA2R contains three TNFR-Cys repeats. It is a new

Target Details

member of the tumor necrosis factor receptor family that has been shown to be highly expressed in ectodermal derivatives during embryonic development and binds to ectodysplasin-A2 (EDA-A2). TNFRSF27 / EDA2R is a receptor for EDA isoform A2, but not for EDA isoform A1. TNFRSF27 / EDA2R mediates the activation of the NF-kappa-B and JNK pathways. The activation seems to be mediated by binding to TRAF3 and TRAF6.

Ectodysplasin, a member of the tumor necrosis factor family, is encoded by the anhidrotic ectodermal dysplasia EDA gene. Mutations in EDA give rise to a clinical syndrome characterized by loss of hair, sweat glands, and teeth. EDA-A1 and EDA-A2 are two isoforms of ectodysplasin that differ only by an insertion of two amino acids. This insertion functions to determine receptor binding specificity, such that EDA-A1 binds only the receptor EDAR, whereas EDA-A2 binds only the related, but distinct, X-linked ectodysplasin-A2 receptor (XEDAR).

Synonym: EDA2R

Molecular Weight:	42.3 kDa
-------------------	----------

UniProt:	D3ZAX4
----------	------------------------

Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling


Format:	Lyophilized
---------	-------------

Reconstitution:	Please refer to the printed manual for detailed information.
-----------------	--

Buffer:	Lyophilized from sterile PBS, pH 7.4
---------	--------------------------------------

Storage:	4 °C, -20 °C, -80 °C
----------	----------------------

Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
------------------	---



Western Blotting

Image 1.