# antibodies.com

# Datasheet for ABIN7321203 LTBR Protein (Fc Tag)

Image



#### Overview

Quantity:	100 µg
Target:	LTBR
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This LTBR protein is labelled with Fc Tag.

### Product Details


Purpose:	Recombinant Rat LTBR/TNFRSF3 Protein (Fc Tag)
Sequence:	Met1-Ala218
Characteristics:	A DNA sequence encoding the rat LTBR (Q5U2S8) (Met1-Ala218) was expressed with the Fc region of human IgG1 at the C-terminus.
Purity:	(9.8+79.7) % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per $\mu g$ of the protein as determined by the LAL method

## Target Details

Target:	LTBR
Alternative Name:	LTBR/TNFRSF3 (LTBR Products)
Background:	Background: LTBR (lymphotoxin beta receptor (TNFR superfamily, member 3)) is a member of the tumor necrosis factor (TNF) family of receptors. Tumor necrosis factor receptor is a
	trimeric cytokine receptor that binds tumor necrosis factors. The receptor cooperates with an

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN7321203 | 09/09/2023 | Copyright antibodies-online. All rights reserved.

	adaptor protein (such as TRADD, TRAF, RIP), which is important in determining the outcome of
	the response. LTBR is expressed on the surface of most cell types, including cells of epithelial
	and myeloid lineages, but not on T and B lymphocytes. LTBR specifically binds the lymphotoxin
	membrane form (a complex of lymphotoxin-alpha and lymphtoxin-beta). LTBR and its ligand
	play a role in the development and organization of lymphoid tissue and tranformed cells.
	Activation of this protein can trigger apoptosis. Not only does the LTBR help trigger apoptosis, it
	can lead to the release of the cytokine interleukin 8. Overexpression of LTBR in HEK293 cells
	increases IL-8 promoter activity and leads to IL-8 release. It is also essential for development
	and organization of the secondary lymphoid organs and chemokine release.
	Synonym: LTBR
Molecular Weight:	48.5 kDa
UniProt:	Q5U2S8
Pathways:	NF-kappaB Signaling
Application Details	
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Please refer to the printed manual for detailed information.
Buffer:	Lyophilized from sterile PBS, pH 7.4
Storage:	4 °C,-20 °C,-80 °C
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted
	samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN7321203 | 09/09/2023 | Copyright antibodies-online. All rights reserved.