

Datasheet for ABIN7321206
CD2 Protein (CD2) (Fc Tag)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µg
Target:	CD2
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This CD2 protein is labelled with Fc Tag.

Product Details

Purpose:	Recombinant Rat CD2 Protein (Fc Tag)
Sequence:	Met1-Pro202
Characteristics:	A DNA sequence encoding the rat CD2 (Q5I0M6) (Met1-Pro202) was expressed, fused with the Fc region of human IgG1 at the C-terminus.
Purity:	> 95 % as determined by SDS-PAGE
Endotoxin Level:	< 1.0 EU per µg of the protein as determined by the LAL method

Target Details

Target:	CD2
Alternative Name:	CD2 (CD2 Products)
Background:	Background: T-cell surface antigen CD2, also known as T-cell surface antigen T11/Leu-5, and SRBC, is a single-pass type I membrane protein. It contains one Ig-like C2-type domain and one Ig-like V-type domain. CD2 is a cell adhesion molecule expressed on T cells and is recognized

Target Details

as a target for CD48 (rats) and CD58 (humans). CD2 has been shown to set quantitative thresholds in T cell activation both in vivo and in vitro. Further, intracellular CD2 signaling pathways and networks are being discovered by the identification of several cytosolic tail binding proteins. CD2 interacts with lymphocyte function-associated antigen (LFA-3) and CD48/BCM1 to mediate adhesion between T-cells and other cell types. CD2 is implicated in the triggering of T-cells, the cytoplasmic domain of CD2 is implicated in the signaling function. The complex of CD2 and CD58 also plays an important role in enhancing the adhesion of T lymphocytes to target cells, and promoting hyperplasia and activation of T lymphocytes. As a cell surface glycoprotein, CD2 expressed on most human T cells and natural killer (NK) cells and plays an important role in mediating cell adhesion in both T-lymphocytes and in signal transduction. Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy

Synonym: CD2

Molecular Weight: 47.5 kDa

UniProt: [Q5I0M6](#)

Application Details

Restrictions: For Research Use only

Handling


Format: Lyophilized

Reconstitution: Please refer to the printed manual for detailed information.

Buffer: Lyophilized from sterile PBS, pH 7.4

Storage: 4 °C, -20 °C, -80 °C

Storage Comment: Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.



Western Blotting

Image 1.