

Datasheet for ABIN7321222

ErbB2/Her2 Protein (AA 4-656) (His tag)

1 Image



Go to Product page

\sim				
()	Ive	r\ /	\cap	Λ.
\cup	$\lor \lor \vdash$	I V I	\Box	٧V

Quantity:	100 μg
Target:	ErbB2/Her2
Protein Characteristics:	AA 4-656
Origin:	Rat
Source:	HEK-293 Cells
Protein Type:	Recombinant
Purification tag / Conjugate:	This ErbB2/Her2 protein is labelled with His tag.

Product Details

Purpose:	Recombinant Rat HER2/ErbB2 Protein (aa 4-656, His Tag)	
Sequence:	Met 4-Thr 656	
Characteristics:	A DNA sequence encoding the rat ErbB2 (AAH61863.1) extracellular domain (Met 4-Thr 656) was expressed, fused with a polyhistidine tag at the C-terminus.	
Purity:	> 95 % as determined by SDS-PAGE	
Endotoxin Level:	$<$ 1.0 EU per μg of the protein as determined by the LAL method	

Target Details

Target:	ErbB2/Her2
Alternative Name:	HER2/ErbB2 (ErbB2/Her2 Products)
Background: Background: Epidermal growth factor receptor 2 (HER2), also known as ErbB2, NEU	

CD340, is a type I membrane glycoprotein, and belongs to the epidermal growth factor (EGF) receptor family. HER2 protein cannot bind growth factors due to the lacking of ligand binding domain of its own and autoinhibited constitutively. However, HER2 forms a heterodimer with other ligand-bound EGF receptor family members, therefore stabilizes ligand binding and enhances kinase-mediated activation of downstream molecules. HER2 plays a key role in development, cell proliferation and differentiation. HER2 gene has been reported to associate with malignancy and a poor prognosis in numerous carcinomas, including breast, prostate, ovarian, lung cancers and so on.Immune Checkpoint Immunotherapy Cancer Immunotherapy Targeted Therapy

Synonym: Receptor tyrosine-protein kinase erbB-2; Epidermal growth factor receptor-related protein; Proto-oncogene Neu; Proto-oncogene c-ErbB-2; p185erbB2; p185neu; CD340; ERBB2;ENV;ENVW;ERVWE1;HER-2;HER-2/neu;HER2;HERV-7q;HERV-W-

ENV;HERV7Q;HERVW;HERVWENV;MLN 19;MLN19

Molecular Weight:

71.3 kDa

Pathways:

RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Skeletal Muscle Fiber Development


Application Details

Restrictions:

For Research Use only

Handling

Format:	Lyophilized	
Reconstitution:	Please refer to the printed manual for detailed information.	
Buffer:	Lyophilized from sterile PBS, pH 7.4	
Storage:	4 °C,-20 °C,-80 °C	
Storage Comment:	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C.	
	Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted	
	samples are stable at < -20°C for 3 months.	



Western Blotting

Image 1.