

Datasheet for ABIN733088

anti-DcR2 antibody**2** Images[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	DcR2 (TNFRSF10D)
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DcR2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DcR2
Isotype:	IgG
Cross-Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

Target:	DcR2 (TNFRSF10D)
Alternative Name:	Dcr2/Tnfrsf10d (TNFRSF10D Products)
Background:	Optional[synonyms]: TNFRSF10D, Decoy receptor 2, CD 264, CD264, CD-264, DcR 2, decoy with truncated death domain, ID, TNF receptor related receptor for TRAIL, TNF related apoptosis inducing ligand receptor 4, TNFRSF 10D, TRAIL R4, TRAIL receptor 4, TRAIL receptor with a truncated death domain, TRAILR4, TRIALR 4, TRUNDD, Tumor necrosis factor receptor

Target Details


	superfamily member 10d decoy with truncated death domain, Tumor necrosis factor receptor superfamily member 10D precursor, TR10D_HUMAN.
Gene ID:	8793
UniProt:	Q9UBN6
Pathways:	Apoptosis

Application Details

Application Notes:	WB: 1:100-1000, FCM: 1:20-100, IHC-P: 1:100-500, IF(IHC-P): 1:50-200 Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



Immunohistochemistry (Paraffin-embedded Sections)

Image 1. Formalin-fixed and paraffin embedded human laryngeal tissue labeled with Rabbit Anti-DcR2/TNFRSF10D Polyclonal Antibody, Unconjugated (ABIN733088) at 1:200 followed by conjugation to the secondary antibody and DAB staining



Flow Cytometry

Image 2. Mouse kidney cells probed with Rabbit Anti-DcR2/TNFRSF10D Polyclonal Antibody, Unconjugated (ABIN733088) (green) at 1:50 for 40 minutes followed by incubation with a FITC conjugated secondary compared to control cells (blue).