

[Go to Product page](#)

Datasheet for ABIN734155
anti-FDCSP antibody (AA 18-85) (Biotin)

Overview

Quantity:	100 µL
Target:	FDCSP (C4orf7)
Binding Specificity:	AA 18-85
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FDCSP antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from rat FDC-SP
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Horse
Purification:	Purified by Protein A.

Target Details

Target:	FDCSP (C4orf7)
Alternative Name:	FDC-SP (C4orf7 Products)

Target Details

Background:	<p>Synonyms: FDC, FDCs, Follicular dendritic cell, Follicular dendritic cell secreted protein, Follicular dendritic cell, Follicular dendritic cell secreted peptide, FDC secreted protein.</p> <p>Background: Follicular dendritic cells are cells that facilitate antigen recognition by B cells in follicles. This kind of dendritic cell is not bone marrow derived and is not a kind of blood cell. It is purely a resident of the follicles of secondary lymphoid organs. B cells form germinal centers around follicular dendritic cells in lymphoid organs. Dendritic cells form from monocytes, white blood cells which circulate in the body and, depending on the right signal, can turn into dendritic cells or macrophages. The monocytes in turn are formed from stem cells in the bone marrow. In normal tissue this antibody identifies dendritic cells and a proportion of B lymphocytes.</p> <p>Evidence in pathological tissue and functional studies suggests it binds to an epitope expressed by antigen presenting cells. Within the macrophage/dendritic cell populations the epitope seen by RFD1 is coexpressed with RFD7 by a subset of cells that exhibit suppressive activity on T cell stimulation.</p>
-------------	--

Gene ID:	494224
----------	--------

UniProt:	Q5R1M6
----------	------------------------

Application Details

Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.

Handling

Expiry Date: 12 months