antibodies - online.com







anti-HMGB4 antibody (AA 31-130)





Overview

Quantity:	100 μL
Target:	HMGB4
Binding Specificity:	AA 31-130
Reactivity:	Human, Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HMGB4 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HMGB4
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Pig
Purification:	Purified by Protein A.

Target Details

Target: HMGB4


Target Details

Target Details		
Alternative Name:	HMGB4 (HMGB4 Products)	
Background:	Synonyms: dJ1007G16.5, High mobility group protein B4, HMGB4	
	Background: High Mobility Group (HMG) proteins, a group of chromosomal proteins common	
	to all eukaryotes, bind DNA in a non-sequence-specific fashion to promote chromatin function	
	and gene regulation, helping with transcription, replication, recombination, and DNA repair.	
	HMGB4 is HMG2-like, isoform 2, high-mobility group box 4. This gene can be found on	
	Chromosome 1. HMGB4 contains two HMG-box regions, and is found in a variety of eukaryotic	
	chromosomal proteins. HMGB proteins are phosphorylated to various extents. The existence o	
	differentially modified forms increases the number of distinct HMGB protein variants in plant	
	chromatin that may be adapted to certain functions.	
Gene ID:	127540	
UniProt:	Q8WW32	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	

Handling


Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

Images




Immunohistochemistry

Image 1. Formalin-fixed and paraffin embedded human colon carcinoma labeled with Anti-HMGB4 Polyclonal Antibody, Unconjugated (ABIN734918) at 1:200 followed by conjugation to the secondary antibody and DAB stainin



ELISA

Image 2. Antigen: $0.2 \mu g/100 \mu L$ Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000; Secondary: HRP conjugated Goat-Anti-Rabbit IgG at 1: 5000; TMB staining; Read the data in MicroplateReader by 450



ELISA

Image 3. Antigen: 0.2ug/100ul, Primary: Antiserum, 1:500, 1:1000, 1:2000, 1:4000, 1:8000, 1:16000, 1:32000, Secondary: HRP conjugated Goat-Anti-Rabbit IgG at 1: 5000, TMB staining, Read the data in MicroplateReader by 450nm