# antibodies - online.com







# anti-IL-11 antibody (AA 22-120) (FITC)



#### Overview

Quantity:	100 μL
Target:	IL-11 (IL11)
Binding Specificity:	AA 22-120
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IL-11 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human IL-11
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Cow,Pig,Guinea Pig
Purification:	Purified by Protein A.

## **Target Details**

Target:	IL-11 (IL11)
Alternative Name:	II-11 (IL11 Products)

## **Target Details**

•	
Background:	Synonyms: Interleukin-11, Adipogenesis inhibitory factor, AG, IL 11, IL11, Interleukin 11,
	Oprelvekin, IL11_HUMAN.
	Background: Interleukin 11 is a pleiotropic cytokine produced by mesenchymal-derived
	adherent cells. IL11 shares many functions of IL6 and LIF, including potentiation of
	megakaryocyte activity, enhancement of human myeloma cell proliferation, and enhancement
	of hepatic acute phase protein production. Interleukin 11 directly stimulates the proliferation of
	hematopoietic stem cells and megakaryocyte progenitor cells, and induces megakaryocyte
	maturation resulting in increased platelet production. Clinically it is used to prevent severe
	thrombocytopenia and the reduction of the need for platelet transfusion following
	myelosuppressive chemotherapy.
Gene ID:	3589
UniProt:	P20809
Pathways:	JAK-STAT Signaling, Negative Regulation of Hormone Secretion
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months