

Datasheet for ABIN735068 anti-IL16 antibody (AA 25-125)



Overview

Target:

IL16

| Quantity: | 100 μL |
|-----------------------|--|
| Target: | IL16 |
| Binding Specificity: | AA 25-125 |
| Reactivity: | Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This IL16 antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunofluorescence (Cultured Cells) (IF (cc)) |
| Product Details | |
| Immunogen: | KLH conjugated synthetic peptide derived from human IL-16 |
| Isotype: | IgG |
| Cross-Reactivity: | Mouse |
| Predicted Reactivity: | Human,Rat |
| Purification: | Purified by Protein A. |
| Target Details | |

Target Details

| Alternative Name: | IL-16 (IL16 Products) | |
|---------------------|--|--|
| Background: | Synonyms: Interleukin-16, HGNC:5980, HsT19289, Interleukin 16 precursor, LCF, Lymphocyte Chemotactic Factor, IL16, Lymphocyte chemoattractant factor, PrIL16, IL16_HUMAN. Background: IL16 is a proinflammatory cytokine that is chemotactic for CD4+ T lymphocytes, monocytes, and eosinophils. IL16 is synthesized as a precursor molecule (pro IL16) of approximately 68 kDa and 631 amino acid residues lacking a signal peptide. The sequence and structure of IL16 is conserved across species. In addition to inducing chemotaxis, IL16 upregulates the IL2 receptor and upregulates HLADR expression. It also inhibits T cell receptor (TCR)/CD3 dependent activation, and suppresses HIV1 replication in vitro. Sources of IL16 include epithelial cells, mast cells, T lymphocytes (CD4+ and CD8+), macrophages, synovial fibroblasts, and eosinophils. | |
| Gene ID: | 3603 | |
| Application Details | | |
| Application Notes: | WB 1:300-5000 ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. | |
| Storage: | 4 °C,-20 °C | |
| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. | |

| | | 1. | |
|----------|-------|-------|---|
| \vdash | land | ling | |
| 1 1 | ıarıu | 11111 | ٠ |
| | | | |

Expiry Date:

12 months