# antibodies -online.com







# anti-IL-20 antibody (AA 51-150) (Cy5)



#### Overview

Quantity:	100 μL
Target:	IL-20 (IL20)
Binding Specificity:	AA 51-150
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IL-20 antibody is conjugated to Cy5
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from mouse IL-20
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human, Mouse
Purification:	Purified by Protein A.

## **Target Details**

Target:	IL-20 (IL20)
Alternative Name:	II-20 (IL20 Products)

## **Target Details**

9	
Background:	Synonyms: Interleukin-20, four alpha helix cytokine, IL10D, Interleukin 20, ZCYTO10,
	IL20_MOUSE, Cytokine Zcyto10.
	Background: IL20 is a cytokine structurally related to interleukin 10 (IL10). This cytokine has
	been shown to transduce its signal through signal transducer and activator of transcription 3
	(STAT3) in keratinocytes. A specific receptor for this cytokine is expressed in skin and
	upregulated dramatically in psoriatic skin, suggesting a role for this protein in epidermal
	function and psoriasis.
Gene ID:	50604
Pathways:	Protein targeting to Nucleus
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months