# antibodies -online.com





Datasheet for ABIN735554

# anti-DNAJC3 antibody (AA 351-450) (Cy7)

# Go to Product page

#### Overview

| Quantity:            | 100 μL  |
|----------------------|---|
| Target:              | DNAJC3  |
| Binding Specificity: | AA 351-450  |
| Reactivity:          | Mouse, Rat  |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This DNAJC3 antibody is conjugated to Cy7   |
| Application:         | Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human p58ipk/DNAJC3 |
|-----------------------|---|
| Isotype:              | IgG   |
| Cross-Reactivity:     | Mouse, Rat  |
| Predicted Reactivity: | Human,Dog,Cow,Pig,Horse   |
| Purification:         | Purified by Protein A.  |

## **Target Details**

| Target:           | DNAJC3                          |
|-------------------|---------------------------------|
| Alternative Name: | P58ipk/Dnajc3 (DNAJC3 Products) |

## **Target Details**

| •                   |  |
|---------------------|--|
| Background:         | Synonyms: DNAJC3, Protein Kinase Inhibitor P58, DnaJ homolog subfamily C member 3, HP58,           |
|                     | Interferon induced, double stranded RNA activated protein kinase inhibitor, P58, p58 ipk,          |
|                     | P58IPK, PRKRI, Protein kinase inhibitor of 58 kDa, Protein kinase inhibitor p58.                   |
|                     | Background: DNAJC3 may act as an inhibitor of both the autophosphorylation of EIF2AK2/PKR          |
|                     | and the ability of EIF2AK2/PKR to catalyze phosphorylation of the EIF2A. It also functions as a    |
|                     | Co-chaperone of HSPA8/HSC70 and is involved in the unfolded protein response (UPR) during          |
|                     | ER stress. It has been reported that this protein stimulates the ATPase activity of                |
|                     | HSPA8/HSC70.DNAJC3 is also involved in ativiral defense - it is activated in response to           |
|                     | infection by influenza virus through the dissociation of DNAJB1. DNAJC3 is widely expressed        |
|                     | and can be found at high levels in the testis and pancreas (expression in cell lines is variable). |
| Gene ID:            | 5611   |
| UniProt:            | Q13217   |
| Pathways:           | ER-Nucleus Signaling   |
| Application Details |  |
| Application Notes:  | IF(IHC-P) 1:50-200   |
|                     | IF(IHC-F) 1:50-200   |
|                     | IF(ICC) 1:50-200   |
| Restrictions:       | For Research Use only  |
| Handling            |  |
| Format:             | Liquid   |
| Concentration:      | 1 μg/μL  |
| Buffer:             | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and        |
|                     | 50 % Glycerol.   |
| Preservative:       | ProClin  |
| Precaution of Use:  | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be                |
|                     | handled by trained staff only.   |
| Storage:            | -20 °C   |
| Storage Comment:    | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.                  |
| Expiry Date:        | 12 months  |
|                     |  |