

[Go to Product page](#)

Datasheet for ABIN735595

**anti-SPDEF/PSE antibody (AA 251-335) (Biotin)**

## Overview

Quantity:	100 µL
Target:	SPDEF/PSE (SPDEF)
Binding Specificity:	AA 251-335
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SPDEF/PSE antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PDEF
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Dog,Cow,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	SPDEF/PSE (SPDEF)
Alternative Name:	PDEF/SPDEF ( <a href="#">SPDEF Products</a> )

## Target Details

Background:	<p>Synonyms: bA375E1.3, PDEF, Prostate derived Ets factor, PROSTATE EPITHELIUM SPECIC ETS TRANSCRIPTION FACTOR, Prostate epithelium-specic Ets transcription factor, Prostate specic Ets, Prostate-derived Ets factor, Prostate-specic Ets, PSE, RP11-375E1__A.3, SAM POINTED DOMAIN CONTAINING ETS TRANSCRIPTION FACTOR, SAM pointed domain-containing Ets transcription factor, Spdef, SPDEF_HUMAN.</p> <p>Background: SPDEF, also known as PDEF is an ETS transcription factor expressed in the prostate epithelial cells. It is thought to act as an androgen-independent transactivator of the prostate-specific antigen (PSA) promoter and also as a transcriptional activator for the SERPINB5 promoter. It is highly expressed in prostate carcinoma cells and is thought to be involved in the regulation of the prostate gland and/or prostate cancer development. It is also expressed in epithelial breast carcinoma.</p>
-------------	---

Gene ID:	25803
----------	-------

## Application Details

Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months