antibodies - online.com





anti-Prostaglandin E Synthase antibody (AA 85-152) (Biotin)



()	۱ ۱	\cap	r	/1	\cap	۱ ۸	1
0	'V	ㄷ	I١	νı	ㄷ	٧	۷

Quantity:	100 μL	
Target:	Prostaglandin E Synthase (PTGES)	
Binding Specificity:	AA 85-152	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Prostaglandin E Synthase antibody is conjugated to Biotin	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), ELISA	
Product Details		

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PTGEs
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Pig,Guinea Pig
Purification:	Purified by Protein A.

Target Details

Target:	Prostaglandin E Synthase (PTGES)	
Alternative Name:	PTGEs (PTGES Products)	
Background:	ground: Synonyms: PGES, MPGES, PIG12, PP102, PP1294, MGST-IV, MGST1L1, TP53I12, mPGES-	

MGST1-L1, Prostaglandin E synthase, Microsomal glutathione S-transferase 1-like 1, Microsomal prostaglandin E synthase 1, p53-induced gene 12 protein, PTGES, MPGES1 Background: Prostaglandin E synthase catalyses conversion of COX derived PGH2 to PGE2. It is a member of the MAPEG family. It may contribute to the pathogenesis of collagen induced arthritis and mediate acute pain during inflammatory responses. The protein encoded by this gene is a glutathione-dependent prostaglandin E synthase. The expression of this gene has been shown to be induced by proinflammatory cytokine interleukin 1 beta (IL1B). Its expression can also be induced by tumor suppressor protein TP53, and may be involved in TP53 induced apoptosis. Knockout studies in mice suggest that this gene may contribute to the pathogenesis of collagen-induced arthritis and mediate acute pain during inflammatory responses.

Gene ID:

9536

UniProt:

014684

Application Details

Application Notes: WB 1:300-5000

IHC-P 1:200-400

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months