antibodies -online.com





anti-RELM alpha antibody (AA 51-111) (Cy7)



| \sim | |
|--------|-------|
| ()\/\ | rview |
| | VICVV |

Background:

| Quantity: | 100 μL |
|----------------------|--|
| Target: | RELM alpha |
| Binding Specificity: | AA 51-111 |
| Reactivity: | Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This RELM alpha antibody is conjugated to Cy7 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)) |
| Product Details | |
| Immunogen: | KLH conjugated synthetic peptide derived from mouse RELMa |
| Isotype: | IgG |
| Cross-Reactivity: | Mouse, Rat |
| Purification: | Purified by Protein A. |
| Target Details | |
| Target: | RELM alpha |
| Alternative Name: | Relma/Relm alpha (RELM alpha Products) |
| | |

Synonyms: HIMF, Xcp2, Fizz1, RELMa, Fizz-1, RELMalpha, RELM-alpha, 18119L16Rik, Resistin-

like alpha, Cysteine-rich secreted protein A12-gamma, Cysteine-rich secreted protein FIZZ1,

Hypoxia-induced mitogenic factor, Parasite-induced macrophage novel gene 1 protein, Retnla, Pmng1

Background: A family of resistin-like molecules (RELMs) has been identified in rodents and humans. RELM alpha belongs to a unique family of tissue-specific cytokines termed FIZZ (found in inflammatory zone) and RELM. The three known members of this family, Resistin, RELM alpha and RELM beta are 85-94 amino acid secreted proteins sharing a conserved C-terminal domain. RELM alpha and Resistin are secreted exclusively by adipocytes while RELM beta is expressed in the epithelium of the colon and small bowel. The RELMs together with resistin comprise a class of tissue-specific signaling molecules. The physiological role and molecular targets of RELM alpha are still unknown.

Gene ID:

57262

UniProt:

Q9EP95

Application Details

Application Notes: IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |