

Datasheet for ABIN736971

anti-CRYGD antibody[Go to Product page](#)

2 Images

1 Publication

Overview

Quantity:	100 µL
Target:	CRYGD
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CRYGD antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide of CCP
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Cross-Reactivity (Details):	CCP
Purification:	Purified by Protein A.

Target Details

Target:	CRYGD
Alternative Name:	Ccp (CRYGD Products)
Target Type:	Chemical
Background:	Synonyms: cyclic citrullinated peptide.

Target Details

Background: CCP

Application Details

Application Notes: WB 1:300-5000
ELISA 1:500-1000
IHC-P 1:200-400

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.


Storage: 4 °C,-20 °C

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date: 12 months


Publications

Product cited in: Sun, Liu, Ma, Wang, Guo, Teng, Jiang: "Effects of forced swimming stress on thyroid function, pituitary thyroid-stimulating hormone and hypothalamus thyrotropin releasing hormone expression in adrenalectomy Wistar rats." in: **Experimental and therapeutic medicine**, Vol. 12, Issue 5, pp. 3167-3174, (2016) ([PubMed](#)).



Immunohistochemistry

Image 1. Formalin-fixed and paraffin embedded human colon carcinoma labeled with Anti-CCP Polyclonal Antibody, Unconjugated (ABIN736971) at 1:200 followed by conjugation to the secondary antibody and DAB staining



Immunohistochemistry (Paraffin-embedded Sections)

Image 2. Formalin-fixed and paraffin embedded human colon carcinoma labeled with Anti-CCP Polyclonal Antibody, Unconjugated at 1:200 followed by conjugation to the secondary antibody and DAB staining