

Datasheet for ABIN737018 anti-HTR2A antibody (AA 65-160) (Biotin)



Overview

Alternative Name:

Background:

Quantity:	100 μL
Target:	HTR2A
Binding Specificity:	AA 65-160
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HTR2A antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human 5-HTR2A
Immunogen:	KLH conjugated synthetic peptide derived from human 5-HTR2A
Isotype:	IgG
Isotype: Cross-Reactivity:	lgG Human, Mouse, Rat

Synonyms: HTR2, 5-HT2A, 5-hydroxytryptamine receptor 2A, 5-HT-2, 5-HT-2A, Serotonin

5-HTR2 (HTR2A Products)

	receptor	2A.	HTR2A
--	----------	-----	-------

Background: G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for various drugs and psychoactive substances, including mescaline, psilocybin, 1-(2,5-dimethoxy-4-iodophenyl)-2-aminopropane (DOI) and lysergic acid diethylamide (LSD). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors. Beta-arrestin family members inhibit signaling via G proteins and mediate activation of alternative signaling pathways. Signaling activates phospholipase C and a phosphatidylinositol-calcium second messenger system that modulates the activity of phosphatidylinositol 3-kinase and promotes the release of Ca(2+) ions from intracellular stores. Affects neural activity, perception, cognition and mood. Plays a role in the regulation of behavior, including responses to anxiogenic situations and psychoactive substances. Plays a role in intestinal smooth muscle contraction, and may play a role in arterial vasoconstriction.

Gene ID:	3356
UniProt:	P28223
Pathways:	JAK-STAT Signaling, Inositol Metabolic Process, Regulation of Carbohydrate Metabolic Process

Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months