



[Go to Product page](#)

Datasheet for ABIN737094
anti-ADRA2A antibody (AA 385-450) (Cy3)

Overview

Quantity:	100 µL
Target:	ADRA2A
Binding Specificity:	AA 385-450
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ADRA2A antibody is conjugated to Cy3
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ADRA2
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Cow,Pig,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	ADRA2A
Alternative Name:	ADRA2 (ADRA2A Products)

Target Details

Background: Synonyms: ADRA2, ADRAR, ZNF32, ADRA2R, ALPHA2AAR, Alpha-2A adrenergic receptor, Alpha-2 adrenergic receptor subtype C1, Alpha-2A adrenoreceptor, Alpha-2A adrenoceptor, Alpha-2AAR, ADRA2A

Background: Alpha-2 adrenergic receptors mediate the catecholamine-induced inhibition of adenylate cyclase through the action of G proteins. The rank order of potency for agonists of this receptor is oxymetazoline > clonidine > epinephrine > norepinephrine > phenylephrine > dopamine > p-syneprine > p-tyramine > serotonin = p-octopamine. For antagonists, the rank order is yohimbine > phentolamine = mianserine > chlorpromazine = spiperone = prazosin > propranolol > alprenolol = pindolol.

Gene ID: 150

UniProt: [P08913](#)

Pathways: [EGFR Signaling Pathway](#), [Negative Regulation of Hormone Secretion](#), [Carbohydrate Homeostasis](#), [cAMP Metabolic Process](#), [Regulation of G-Protein Coupled Receptor Protein Signaling](#), [Negative Regulation of Transporter Activity](#)

Application Details

Application Notes: IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months