antibodies -online.com





anti-SLC39A7 antibody (AA 201-300)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL	
Target:	SLC39A7	
Binding Specificity:	AA 201-300	
Reactivity:	Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This SLC39A7 antibody is un-conjugated	
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human SLC39A7
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Dog,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

|--|

Target Details

rarget betails		
Alternative Name:	SLC39A7 (SLC39A7 Products)	
Background:	Synonyms: D6S115E, D6S2244E, H2 KE4, Histidine rich membrane protein Ke4, Histidine-rich	
	membrane protein Ke4, HKE4, HLA class II region expressed gene KE4, KE4, Ke4 gene mouse,	
	human homolog of, Really interesting new gene 5 protein, RING5, S39A7_HUMAN, SLC39A7,	
	Solute carrier family 39 zinc transporter, member 7, Solute carrier family 39 member 7, solute	
	carrier family 39 zinc transporter member 7, Zinc transporter SLC39A7, ZIP7.	
	Background: Zinc is an essential cofactor for more than 50 classes of enzymes and is involved	
	in protein, nucleic acid, carbohydrate, and lipid metabolism, plus the control of gene	
	transcription, growth, development, and differentiation. Zinc cannot passively diffuse across	
	cell membranes and requires specific transporters, such as SLC39A7, to enter the cytosol from	
	both the extracellular environment and from intracellular storage compartments.	
Gene ID:	7922	
Pathways:	Transition Metal Ion Homeostasis	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	

Handling

Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months