antibodies - online.com







anti-CKLF antibody (AA 1-100)





Overview

Quantity:	100 μL
Target:	CKLF
Binding Specificity:	AA 1-100
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CKLF antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CKLF
Isotype:	IgG
Cross-Reactivity:	Human, Mouse, Rat
Predicted Reactivity:	Rabbit
Purification:	Purified by Protein A.

Target Details

|--|

Target Details


Alternative Name:	CKLF (CKLF Products)
Background:	Synonyms: C32, CKLF1, CKLF2, CKLF3, CKLF4, UCK-1, HSPC224, Chemokine-like factor, CKLF, UNQ410/PR0772
	Background: May play an important role in inflammation and regeneration of skeletal muscle.
	Partly inhibited by interleukin 10. Isoform 1 has chemotactic response in rat monocytes,
	neutrophils, and lymphocytes.
Gene ID:	51192
UniProt:	Q9UBR5

Application Details

Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months




Immunohistochemistry

Image 1. Formalin-fixed and paraffin embedded human liver carcinoma labeled with Anti-CKLF Polyclonal Antibody, Unconjugated (ABIN739650) at 1:200, followed by conjugation to the secondary antibody and DAB staining



Immunohistochemistry

Image 2. Formalin-fixed and paraffin embedded mouse spleen labeled with Anti-CKLF Polyclonal Antibody, Unconjugated (ABIN739650) at 1:200, followed by conjugation to the secondary antibody and DAB staining



Immunohistochemistry

Image 3. Formalin-fixed and paraffin embedded rat liver labeled with Anti-CKLF Polyclonal Antibody, Unconjugated (ABIN739650) at 1:200, followed by conjugation to the secondary antibody and DAB staining