

Datasheet for ABIN739980
anti-NCR3 antibody (AA 81-180)[Go to Product page](#)

1 Image

Overview

Quantity:	100 µL
Target:	NCR3
Binding Specificity:	AA 81-180
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NCR3 antibody is un-conjugated
Application:	ELISA, Flow Cytometry (FACS), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NCR3
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Rat,Dog,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NCR3
Alternative Name:	NCR3 (NCR3 Products)
Background:	Synonyms: 1C7, MALS, CD337, LY117, NKp30, Natural cytotoxicity triggering receptor 3,

Target Details

Activating natural killer receptor p30, Natural killer cell p30-related protein, NK-p30, NCR3

Background: Cytotoxicity-activating receptor that contributes to the increased efficiency of activated natural killer (NK) cells to mediate tumor cell lysis. Engagement of NCR3 by BAG6 also promotes dendritic cell (DC) maturation, both through killing those DCs that did not properly acquire a mature phenotype, and inducing NK cells to release TNFA and IFNG, which promotes DC maturation.


Gene ID:	259197
UniProt:	O14931
Pathways:	Regulation of Leukocyte Mediated Immunity , Positive Regulation of Immune Effector Process

Application Details

Application Notes:	ELISA 1:500-1000 FCM 1:20-100 IHC-P 1:200-400
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months



Immunohistochemistry

Image 1. Formalin-fixed and paraffin embedded human lung carcinoma labeled with Anti-NCR3 Polyclonal Antibody, Unconjugated (ABIN739980) at 1:200 followed by conjugation to the secondary antibody and DAB staining