## antibodies -online.com





## anti-NCR3 antibody (AA 81-180) (Biotin)



| iew |
|-----|
|     |

| Overview              |   |
|-----------------------|---|
| Quantity:             | 100 μL  |
| Target:               | NCR3  |
| Binding Specificity:  | AA 81-180   |
| Reactivity:           | Human   |
| Host:                 | Rabbit  |
| Clonality:            | Polyclonal  |
| Conjugate:            | This NCR3 antibody is conjugated to Biotin  |
| Application:          | ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))                    |
| Product Details       |   |
| Immunogen:            | KLH conjugated synthetic peptide derived from human NCR3                              |
| Isotype:              | IgG   |
| Cross-Reactivity:     | Human   |
| Predicted Reactivity: | Rat,Dog,Horse,Rabbit  |
| Purification:         | Purified by Protein A.  |
| Target Details        |   |
| Target:               | NCR3  |
| Alternative Name:     | NCR3 (NCR3 Products)  |
| Background:           | Synonyms: 1C7, MALS, CD337, LY117, NKp30, Natural cytotoxicity triggering receptor 3, |

Activating natural killer receptor p30, Natural killer cell p30-related protein, NK-p30, NCR3 Background: Cytotoxicity-activating receptor that contributes to the increased efficiency of activated natural killer (NK) cells to mediate tumor cell lysis. Engagement of NCR3 by BAG6 also promotes dendritic cell (DC) maturation, both through killing those DCs that did not properly acquire a mature phenotype, and inducing NK cells to release TNFA and IFNG, which promotes DC maturation.

Gene ID: 259197

UniProt: 014931

Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process Pathways:

**Application Details** 

IHC-P 1:200-400 **Application Notes:** 

Restrictions: For Research Use only

Handling

Format: Liquid Concentration:  $1 \mu g/\mu L$ Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

handled by trained staff only.

Storage: -20 °C

Store at -20°C for 12 months Storage Comment:

**Expiry Date:** 12 months