

Datasheet for ABIN740019

anti-NKp44/NCR2 antibody (AA 201-276) (HRP)



Go to Product page

| \sim | | |
|---------|-------|------------|
| Ove | ·r\/I | ΘW |
| \circ | | C * * |

Background:

| Quantity: | 100 μL |
|---|--|
| Target: | NKp44/NCR2 (NCR2) |
| Binding Specificity: | AA 201-276 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This NKp44/NCR2 antibody is conjugated to HRP |
| Application: | ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |
| | |
| Product Details | |
| Product Details Immunogen: | KLH conjugated synthetic peptide derived from human CD336/NCR2 |
| | KLH conjugated synthetic peptide derived from human CD336/NCR2 |
| Immunogen: | |
| Immunogen: Isotype: | IgG |
| Immunogen: Isotype: Cross-Reactivity: | IgG Human |
| Immunogen: Isotype: Cross-Reactivity: Purification: | IgG Human |

Synonyms: LY95, CD336, NKP44, NK-p44, dJ149M18.1, Natural cytotoxicity triggering receptor

Target Details

Expiry Date:

12 months

| rarget Details | | |
|---------------------|---|--|
| | 2, Lymphocyte antigen 95 homolog, NK cell-activating receptor, Natural killer cell p44-related protein, NCR2 | |
| | Background: Cytotoxicity-activating receptor that may contribute to the increased efficiency of | |
| | activated natural killer (NK) cells to mediate tumor cell lysis. | |
| Gene ID: | 9436 | |
| UniProt: | 095944 | |
| Application Details | | |
| Application Notes: | IHC-P 1:200-400 | |
| | IHC-F 1:100-500 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should b handled by trained staff only. | |
| Handling Advice: | Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| | | |