



[Go to Product page](#)

Datasheet for ABIN740357
anti-CYP21A2 antibody (AA 101-200) (Biotin)

Overview

Quantity:	100 µL
Target:	CYP21A2
Binding Specificity:	AA 101-200
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CYP21A2 antibody is conjugated to Biotin
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CYP21
Isotype:	IgG
Cross-Reactivity:	Rat
Predicted Reactivity:	Human,Mouse,Dog,Cow,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target:	CYP21A2
Alternative Name:	CYP21 (CYP21A2 Products)

Target Details

Background: Synonyms: Cytochrome P450 21, cytochrome P450, family 21, subfamily A, polypeptide 2, 21-OH, 21OHA, 21OHB, CA21H, CAH1, CHOLESTEROL 21-HYDROXYLASE, CPS1, CYP21, Cyp21-ps1, Cyp21a1, CYP21A2, Cyp21a2-ps, CYP21B, CYP21OH-A, MGC150536, MGC150537, MGC156449, Oh21-1, Oh21-2, P450C21, P450c21B, CYP21A2, CP21A_HUMAN.

Background: This gene encodes a member of the cytochrome P450 superfamily of enzymes. The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes to the endoplasmic reticulum and hydroxylates steroids at the 21 position. Its activity is required for the synthesis of steroid hormones including cortisol and aldosterone. Mutations in this gene cause congenital adrenal hyperplasia. A related pseudogene is located near this gene, gene conversion events involving the functional gene and the pseudogene are thought to account for many cases of steroid 21-hydroxylase deficiency. Two transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq].

Gene ID: 1589

Pathways: [Metabolism of Steroid Hormones and Vitamin D](#), [Steroid Hormone Biosynthesis](#), [C21-Steroid Hormone Metabolic Process](#)

Application Details

Application Notes: IHC-P 1:200-400
IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C for 12 months.

Handling

Expiry Date: 12 months