# antibodies -online.com







# anti-CCL19 antibody (AA 26-108) (Biotin)



| ( ) | 1/0 | r\ /1 | 014 |   |
|-----|-----|-------|-----|---|
| ( ) | ve  | I V I | -v  | V |
|     |     |       |     |   |

| Quantity:            | 100 μL  |
|----------------------|---|
| Target:              | CCL19   |
| Binding Specificity: | AA 26-108   |
| Reactivity:          | Human, Rat  |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This CCL19 antibody is conjugated to Biotin   |
| Application:         | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from mouse CCL19 |
|-----------------------|---|
| Isotype:              | IgG   |
| Cross-Reactivity:     | Human, Rat  |
| Predicted Reactivity: | Mouse,Pig   |
| Purification:         | Purified by Protein A.                                    |

## **Target Details**

| Target:           | CCL19                             |  |
|-------------------|-----------------------------------|--|
| Alternative Name: | CCL19/MIP-3 beta (CCL19 Products) |  |

## **Target Details**

| Synonyms: ELC, CKb11, MIP3B, Scya19, exodus-3, C-C motif chemokine 19, Epstein-Barr virus-induced molecule 1 ligand chemokine, EBI1 ligand chemokine, Small-inducible cytokine A19, |  |
|---|--|
| Ccl19   |  |
| Background: Strongly chemotactic for naive (L-selectinhi) CD4 T-cells and for CD8 T-cells and   |  |
| weakly attractive for resting B-cells and memory (L-selectinlo) CD4 T-cells. May play a role in   |  |
| promoting encounters between recirculating T-cells and dendritic cells and in the migration of  |  |
| activated B-cells into the T-zone of secondary lymphoid tissues. Binds to chemokine receptor  |  |
| CCR7. Binds to atypical chemokine receptor ACKR4 and mediates the recruitment of beta-  |  |
| arrestin (ARRB1/2) to ACKR4.  |  |
| 24047   |  |
| 070460  |  |
| Positive Regulation of Immune Effector Process  |  |
|   |  |
| WB 1:300-5000   |  |
| IHC-P 1:200-400   |  |
| IHC-F 1:100-500   |  |
| For Research Use only   |  |
|   |  |
| Liquid  |  |
| 1 μg/μL   |  |
| Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and   |  |
| 50 % Glycerol.  |  |
| ProClin   |  |
| This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be   |  |
| handled by trained staff only.  |  |
| -20 °C  |  |
| Store at -20°C for 12 months.   |  |
| 12 months   |  |
|   |  |