

Datasheet for ABIN740559

anti-HVEM antibody (AA 51-150) (HRP)[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	HVEM (TNFRSF14)
Binding Specificity:	AA 51-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HVEM antibody is conjugated to HRP
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TNFRSF14
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Dog,Horse
Purification:	Purified by Protein A.

Target Details

Target:	HVEM (TNFRSF14)
Alternative Name:	TNFRSF14/HVEM (TNFRSF14 Products)

Target Details

Target Type:	Viral Protein
Background:	Synonyms: Tumor necrosis factor receptor superfamily member 14, HVEM, ATAR, CD258, CD258 antigen, CD40 like protein precursor, Herpesvirus entry mediator A, Herpesvirus entry mediator, Herpesvirus entry mediator ligand, HVEA, HVEM, HVEM L, HVEM, LIGHT, LIGHTR, TNFSF 14, TR2, Tumor necrosis factor receptor like gene2, TNFR14_HUMAN. Background: TNFRSF14 is a type I membrane protein belonging to the TNF receptor superfamily. This receptor mediates herpes virus entry into cells during infection. TNFRSF14 is able to inhibit the proliferation, activation, and cytokine production of T cells. It has an extracellular domain containing several cysteine-rich repeats and a short cytoplasmic region containing a TRAF (TNF receptor-associated factor) interaction domain. The extracellular domain of TNFRSF14 interacts with the herpes simplex virus envelope glycoprotein D. TNFRSF14 binds two cellular ligands: lymphotoxin alpha and LIGHT. LIGHT is a transmembrane protein expressed and shed from the surface of activated T cells, exhibits inducible expression, and competes with HSV glycoprotein D for HVEM, a receptor expressed by T lymphocytes. The LIGHT:TNFRSF14 interaction controls immune response functions by cell death induction as well as cell activation. TNFRSF14 is expressed by peripheral blood T cells, B cells, monocytes and in various tissues enriched in lymphoid cells.
Gene ID:	8764
Pathways:	Production of Molecular Mediator of Immune Response , Cancer Immune Checkpoints

Application Details

Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

handled by trained staff only.

Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
------------------	--

Storage:	-20 °C
----------	--------

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
------------------	---

Expiry Date:	12 months
--------------	-----------
