antibodies -online.com





anti-CCL9 antibody (AA 22-122) (FITC)



Go to Product page

\sim			
	N/P	r\/	i⊢₩

Predicted Reactivity:

Purification:

Rat

Purified by Protein A.

Quantity:	100 μL	
Target:	CCL9 (Ccl9)	
Binding Specificity:	AA 22-122	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CCL9 antibody is conjugated to FITC	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from rat CCL9/Macrophage Inflammatory Protein 1 gamma	
Isotype:	IgG	
Specificity:	This antibody may exhibit a slight secondary cross-reactivity with Cerebellar degeneration-related protein 2-like due to a 76 % non-contiguous sequence similarity in the immunogen sequence range.	
Cross-Reactivity:	Mouse	

Target Details

Target:	CCL9 (Ccl9)		
Alternative Name:	Macrophage Inflammatory Protein 1 gamma (Ccl9 Products)		
Background:	Synonyms: MIP-1 gamma, Macrophage inflammatory protein 1-gamma, Macrophage		
	Inflammatory Protein 1 gamma, CCF18, CCL10, CCL9, Chemokine CC mot ligand 9, mMIP1g,		
	SCYA10, SCYA9, Small inducible cytokine A9,MIP 1 gamma, MIP1 gamma, Macrophage		
	Inflammatory Protein 1 gamma.		
	Background: This chemotactic factor that attracts monocytes, lymphocytes, basophils and		
	eosinophils may play a role in neoplasia and inflammatory host responses and can bind		
	heparin. Its highest expression is found in the small intestine and peripheral blood cells. It is		
	induced by interferon gamma, mitogens and interleukin 1 and belongs to the intercrine beta		
	(chemokine CC) family.		
Gene ID:	360579		
Application Details			
Application Notes:	IF(IHC-P) 1:50-200		
	IF(IHC-F) 1:50-200		
	IF(ICC) 1:50-200		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		
Expiry Date:	12 months		