antibodies -online.com





anti-CCL16 antibody (AA 24-120) (Cy5)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL	
Target:	CCL16	
Binding Specificity:	AA 24-120	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CCL16 antibody is conjugated to Cy5	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human CCL16	
Isotype:	IgG	
Predicted Reactivity:	Human	
Purification:	Purified by Protein A.	

Target Details

Target:	CCL16
Alternative Name:	Ccl16/Hcc-4 (CCL16 Products)
Background: Synonyms: C-C mot chemokine 16, chemokine C-C mot ligand 16, CCL16, CKb12, HCC4, IL	

Expiry Date:

Storage Comment:

12 months

l arget Details	
	inducible chemokine, ILINCK, LCC1, Liver CC chemokine 1 precursor, Liver expressed
	chemokine, LMC, Monotactin 1, Mtn1, NCC4, New CC chemokine 4, SCYA16, SCYL4, Small
	inducible cytokine A16 precursor, Small inducible cytokine subfamily A Cys Cys member 16,
	Small-inducible cytokine A16. CCL16_HUMAN
	Background: LEC (liver expressed cytokine) displays chemotactic activity for lymphocytes and
	monocytes but not for neutrophils. It also shows a potent myelosuppressive activity and
	suppresses proliferation of myeloid progenitor cells. Its expression is upregulated by IL10.
Gene ID:	6360
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.