## antibodies -online.com





## anti-CCL16 antibody (AA 24-120) (Cy5.5)



( )	ve	K\ /		A .
	$\cup$	1 V/	-	V۷

Quantity:	100 μL	
Target:	CCL16	
Binding Specificity:	AA 24-120	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CCL16 antibody is conjugated to Cy5.5	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CCL16
Isotype:	IgG
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

## **Target Details**

Target:	CCL16
Alternative Name:	Ccl16/Hcc-4 (CCL16 Products)
Background: Synonyms: C-C mot chemokine 16, chemokine C-C mot ligand 16, CCL16, CKb12, HCC-	

Storage:

Expiry Date:

Storage Comment:

rarget Details		
	inducible chemokine, ILINCK, LCC1, Liver CC chemokine 1 precursor, Liver expressed	
	chemokine, LMC, Monotactin 1, Mtn1, NCC4, New CC chemokine 4, SCYA16, SCYL4, Small	
	inducible cytokine A16 precursor, Small inducible cytokine subfamily A Cys Cys member 16,	
	Small-inducible cytokine A16. CCL16_HUMAN	
	Background: LEC (liver expressed cytokine) displays chemotactic activity for lymphocytes and	
	monocytes but not for neutrophils. It also shows a potent myelosuppressive activity and	
	suppresses proliferation of myeloid progenitor cells. Its expression is upregulated by IL10.	
Gene ID:	6360	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

-20 °C

12 months