antibodies -online.com





anti-CLEC1A antibody (AA 151-280) (Cy3)



Go to Product page

| \sim | | | |
|--------|-----|-----|-----|
| | N/P | r\/ | i⊢₩ |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | CLEC1A |
| Binding Specificity: | AA 151-280 |
| Reactivity: | Mouse, Rat |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This CLEC1A antibody is conjugated to Cy3 |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human CLEC1/CLEC1A |
|-----------------------|--|
| Isotype: | IgG |
| Cross-Reactivity: | Mouse, Rat |
| Predicted Reactivity: | Human,Dog,Cow,Pig,Horse,Rabbit,Guinea Pig |
| Purification: | Purified by Protein A. |

Target Details

| Target: | CLEC1A |
|-------------------|--------------------------------|
| Alternative Name: | CLEC1/CLEC1A (CLEC1A Products) |

Target Details

| Background: | Synonyms: CLEC1, CLEC-1, C-type lectin domain family 1 member A, C-type lectin-like receptor | |
|---------------------|--|--|
| | 1, CLEC1A, UNQ569/PR01131 | |
| | Background: This gene encodes a member of the C-type lectin/C-type lectin-like domain | |
| | (CTL/CTLD) superfamily. Members of this family share a common protein fold and have | |
| | diverse functions, such as cell adhesion, cell-cell signalling, glycoprotein turnover, and roles in | |
| | inflammation and immune response. The encoded protein may play a role in regulating | |
| | dendritic cell function. Alternative splice variants have been described but their biological nature | |
| | has not been determined. This gene is closely linked to other CTL/CTLD superfamily members | |
| | on chromosome 12p13 in the natural killer gene complex region. [provided by RefSeq] | |
| Gene ID: | 51267 | |
| UniProt: | Q8NC01 | |
| Application Details | | |
| Application Notes: | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |
| | | |