antibodies -online.com





anti-IGSF8 antibody (AA 101-200)



_							
0	V	е	r١	/	е	٧	V

Quantity:	100 μL
Target:	IGSF8
Binding Specificity:	AA 101-200
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IGSF8 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Flow Cytometry (FACS), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from mouse IGSF8
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Cow,Pig,Horse
Purification:	Purified by Protein A.

Target Details

Target: IGSF8			
	Target:	IGSF8	

Target Details

- Target Details		
Alternative Name:	IGSF8/CD316 (IGSF8 Products)	
Background:	Synonyms: PGRL, EWI-2, KCT-4, ESTM34, R75384, AA033172, AU041109, Immunoglobulin	
	superfamily member 8, IgSF8, CD81 partner 3, Glu-Trp-lle EWI motif-containing protein 2,	
	Keratinocyte-associated transmembrane protein 4, Prostaglandin regulatory-like protein, CD316, Ewi2, Kct4	
	Background: May play a key role in diverse functions ascribed to CD81 and CD9 such as	
	oocytes fertilization or hepatitis C virus function. May regulate proliferation and differentiation	
	of keratinocytes. May be a negative regulator of cell motility: suppresses T-cell mobility	
	coordinately with CD81, associates with CD82 to suppress prostate cancer cell migration,	
	regulates epidermoid cell reaggregation and motility on laminin-5 with CD9 and CD81 as key	
	linkers. May also play a role on integrin-dependent morphology and motility functions. May	
	participate in the regulation of neurite outgrowth and maintenance of the neural network in th	
	adult brain.	
Gene ID:	140559	
UniProt:	Q8R366	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	FCM 1:20-100	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	

Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months