

## Datasheet for ABIN742472 anti-CYP1A2 antibody (AA 221-320) (Biotin)



Go to Product page

Overview	
Quantity:	100 μL
Target:	CYP1A2
Binding Specificity:	AA 221-320
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CYP1A2 antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)),
	Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Product Details Immunogen:	KLH conjugated synthetic peptide derived from human Cytochrome P450 1A2
	KLH conjugated synthetic peptide derived from human Cytochrome P450 1A2
Immunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype: Cross-Reactivity:	IgG Human, Mouse, Rat
Immunogen: Isotype: Cross-Reactivity: Purification:	IgG Human, Mouse, Rat
Immunogen: Isotype: Cross-Reactivity: Purification: Target Details	IgG Human, Mouse, Rat Purified by Protein A.

Cytochrome P450 4, Cytochrome P450-P3, CYP1A2

Background: Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver microsomes, this enzyme is involved in an NADPH-dependent electron transport pathway. It oxidizes a variety of structurally unrelated compounds, including steroids, fatty acids, and xenobiotics. Most active in catalyzing 2-hydroxylation. Caffeine is metabolized primarily by cytochrome CYP1A2 in the liver through an initial N3-demethylation. Also acts in the metabolism of aflatoxin B1 and acetaminophen. Participates in the bioactivation of carcinogenic aromatic and heterocyclic amines. Catalizes the N-hydroxylation of heterocyclic amines and the O-deethylation of phenacetin.

Gene ID: 1544

UniProt: P05177

## **Application Details**

Application Notes: WB 1:300-5000

IHC-P 1:200-400 IHC-F 1:100-500

Restrictions: For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C for 12 months.
Expiry Date:	12 months