antibodies - online.com







anti-CXCL10 antibody (AA 22-98)





\sim				
	$ V \cap$	r\/I	19	٨

Quantity:	100 μL	
·	<u> </u>	
Target:	CXCL10	
Binding Specificity:	AA 22-98	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This CXCL10 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP),	
	Immunocytochemistry (ICC)	

Product Details

Purpose:	Monoclonal Antibody to Interferon Gamma Induced Protein 10kDa (IP10)	
Immunogen:	Recombinant Interferon Gamma Induced Protein 10kDa (IP10) corresdonding to Val22~Pro98 with N-terminal His Tag	
Clone:	C3	
Isotype:	IgG1 kappa	
Specificity:	The antibody is a mouse monoclonal antibody raised against IP10. It has been selected for its ability to recognize IP10 in immunohistochemical staining and western blotting.	
Purification:	Protein A + Protein G affinity chromatography	

Target Details

CXCL10	
CXCL10	
Interferon Gamma Induced Protein 10kDa (CXCL10 Products)	
CXCL10, C7, IFI10, INP10, SCYB10, Crg2, GIP10, Mob1, Chemokine C-X-C Motif Ligand 10,	
Small Inducible Cytokine Subfamily B(Cys-X-Cys)Member 10	
Western blotting: 0.5-2 μg/mL	
Immunohistochemistry: 5-20 μg/mL	
Immunocytochemistry: 5-20 μg/mL	
Optimal working dilutions must be determined by end user.	
The thermal stability is described by the loss rate. The loss rate was determined by accelerated	
thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	
degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
date under appropriate storage condition.	
For Research Use only	
Liquid	
PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	
Sodium azide	
This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which	
should be handled by trained staff only.	
4 °C,-20 °C	
Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without	
detectable loss of activity. Avoid repeated freeze-thaw cycles.	
24 months	



Western Blotting

Image 1. Detection of IP10 in Human Placenta lysate using Monoclonal Antibody to Interferon Gamma Induced Protein 10 kDa (IP10)