# antibodies -online.com





## anti-CCL18 antibody (AA 21-89)





Go to Product page

#### Overview


Quantity:	100 μL
Target:	CCL18
Binding Specificity:	AA 21-89
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CCL18 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

#### **Product Details**

Purpose:	Monoclonal Antibody to Pulmonary Activation Regulated Chemokine (PARC)
Immunogen:	Recombinant Pulmonary Activation Regulated Chemokine (PARC) corresdonding to Ala21~Ala89 with N-terminal His Tag
Clone:	C3
Isotype:	IgG2a kappa
Specificity:	The antibody is a mouse monoclonal antibody raised against PARC. It has been selected for its ability to recognize PARC in immunohistochemical staining and western blotting.
Purification:	Protein A + Protein G affinity chromatography

### **Target Details**

Target:	CCL18
Alternative Name:	Pulmonary Activation Regulated Chemokine (CCL18 Products)
Background:	CCL18, AMAC1, SCYA18, DCCK1, MIP-4, CKb7, Alternative Macrophage Activation Associated
	CC Chemokine 1, Chemokine C-C-Motif Ligand 18, Macrophage inflammatory protein 4
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL
	Immunohistochemistry: 5-20 μg/mL
	Immunocytochemistry: 5-20 μg/mL
	Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



#### **Western Blotting**

**Image 1.** Detection of Recombinant PARC, Human using Monoclonal Antibody to Pulmonary Activation Regulated Chemokine (PARC)