

Datasheet for ABIN7426562
anti-Stanniocalcin 1 antibody (AA 28-247)[Go to Product page](#)

2 Images

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | Stanniocalcin 1 (STC1) |
| Binding Specificity: | AA 28-247 |
| Reactivity: | Human |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This Stanniocalcin 1 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details

| | |
|-------------------|---|
| Purpose: | Monoclonal Antibody to Stanniocalcin 1 (STC1) |
| Immunogen: | Recombinant Stanniocalcin 1 (STC1) corresponding to Ser28~Ala247 with N-terminal His and GST Tag |
| Clone: | C13 |
| Isotype: | IgG2a kappa |
| Specificity: | The antibody is a mouse monoclonal antibody raised against STC1. It has been selected for its ability to recognize STC1 in immunohistochemical staining and western blotting. |
| Cross-Reactivity: | Mouse, Pig, Rat |
| Purification: | Protein A + Protein G affinity chromatography |

Target Details


| | |
|-------------------|---|
| Target: | Stanniocalcin 1 (STC1) |
| Alternative Name: | Stanniocalcin 1 (STC1 Products) |
| Pathways: | Hormone Activity |

Application Details

| | |
|--------------------|---|
| Application Notes: | Western blotting: 0.5-3 µg/mL Immunohistochemistry: 5-30 µg/mL Immunocytochemistry: 5-30 µg/mL Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions: | For Research Use only |

Handling

| | |
|--------------------|---|
| Format: | Liquid |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. |
| Expiry Date: | 24 months |



Western Blotting

Image 1. Detection of STC1 in Porcine Testis lysate using Monoclonal Antibody to Stanniocalcin 1 (STC1)

Western Blotting

Image 2. Detection of Recombinant STC1, Human using Monoclonal Antibody to Stanniocalcin 1 (STC1)