

Datasheet for ABIN7426814 anti-TNF alpha antibody (AA 80-235)

2 Images



Go to Product page

()	11		r 1	ri	\sim	1 A /
U	V	Ч	rv	ı	С	V۷

Quantity:	100 μL
Target:	TNF alpha
Binding Specificity:	AA 80-235
Reactivity:	Rabbit
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TNF alpha antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)


Product Details

Purpose:	Monoclonal Antibody to Tumor Necrosis Factor Alpha (TNFa)	
Immunogen:	Recombinant Tumor Necrosis Factor Alpha (TNFa) corresdonding to Leu80~Leu235 (Accession # P04924)	
Clone:	D3	
Isotype:	IgG1 kappa	
Specificity:	The antibody is a mouse monoclonal antibody raised against TNFa. It has been selected for its ability to recognize TNFa in immunohistochemical staining and western blotting.	
Purification:	Protein A + Protein G affinity chromatography	

Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, Positive Regulation of Endopeptidase Activity, Hepatitis C, Protein targeting to Nucleus, Inflammasome Application Details Application Notes: Western blotting: 0.01-3 µg/mL,Immunohistochemistry: 5-30 µg/mL,Immunocytochemistry: 30 µg/mL,Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerat thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious	Target Details		
Background: DIF, TNF-A, TNFSF2, Cachectin, Tumor Necrosis Factor Ligand Superfamily Member 2 UniProt: P04924 Pathways: NF-kappaB Signaling, Apoptosis, Caspase Cascade in Apoptosis, TLR Signaling, Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positi Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, Positive Regulation of Endopeptidase Activity, Hepatitis C, Protein targeting to Nucleus, Inflammasome Application Details Application Notes: Western blotting: 0.01-3 μg/mL,Immunohistochemistry: 5-30 μg/mL,Immunocytochemistry: 30 μg/mL,Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerat thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	Target:	TNF alpha	
UniProt: P04924 Pathways: NF-kappaB Signaling, Apoptosis, Caspase Cascade in Apoptosis, TLR Signaling, Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positi Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, Positive Regulation of Endopeptidase Activity, Hepatitis C, Protein targeting to Nucleus, Inflammasome Application Details Application Notes: Western blotting: 0.01-3 μg/mL,Immunohistochemistry: 5-30 μg/mL,Immunocytochemistry: 30 μg/mL,Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerat thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	Alternative Name:	TNFa (TNF alpha Products)	
Pathways: NF-kappaB Signaling, Apoptosis, Caspase Cascade in Apoptosis, TLR Signaling, Cellular Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positi Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, Positive Regulation of Endopeptidase Activity, Hepatitis C, Protein targeting to Nucleus, Inflammasome Application Details Application Notes: Western blotting: 0.01-3 μg/mL,Immunohistochemistry: 5-30 μg/mL,Immunocytochemistry: 30 μg/mL,Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerate thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	Background:	DIF, TNF-A, TNFSF2, Cachectin, Tumor Necrosis Factor Ligand Superfamily Member 2	
Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, Positive Regulation of Endopeptidase Activity, Hepatitis C, Protein targeting to Nucleus, Inflammasome Application Details Application Notes: Western blotting: 0.01-3 µg/mL,Immunohistochemistry: 5-30 µg/mL,Immunocytochemistry: 30 µg/mL,Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerate thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	UniProt:	P04924	
Application Notes: Western blotting: 0.01-3 μg/mL,Immunohistochemistry: 5-30 μg/mL,Immunocytochemistry: 30 μg/mL,Optimal working dilutions must be determined by end user. Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerate thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	Pathways:	Response to Molecule of Bacterial Origin, Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process, Production of Molecular Mediator of Immune Response, Positive Regulation of Endopeptidase Activity, Hepatitis C, Protein targeting to	
30 μg/mL,Optimal working dilutions must be determined by end user. The thermal stability is described by the loss rate. The loss rate was determined by accelerat thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	Application Details		
thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	Application Notes:	Western blotting: 0.01-3 μg/mL,Immunohistochemistry: 5-30 μg/mL,Immunocytochemistry: 5-30 μg/mL,Optimal working dilutions must be determined by end user.	
Restrictions: For Research Use only	Comment:	degradation and precipitation were observed. The loss rate is less than 5% within the expiration	
	Restrictions:	For Research Use only	


Handling

Format:	Liquid
Concentration:	0.87 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



Immunohistochemistry

Image 1. Detection of TNFa in Rabbit Liver Tissue using Monoclonal Antibody to Tumor Necrosis Factor Alpha (TNFa)



Western Blotting

Image 2. Detection of Recombinant TNFa, Rabbit using Monoclonal Antibody to Tumor Necrosis Factor Alpha (TNFa)