

Datasheet for ABIN7426835  
**anti-Tissue factor antibody (AA 34-251)**[Go to Product page](#)

## 3 Images

## Overview

Quantity:	100 µL
Target:	Tissue factor (F3)
Binding Specificity:	AA 34-251
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Tissue factor antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

## Product Details

Purpose:	Monoclonal Antibody to Tissue Factor (TF)
Immunogen:	Recombinant Tissue Factor (TF) corresponding to Gly34~glu251 with N-terminal His Tag
Clone:	C11
Isotype:	IgG2a kappa
Specificity:	The antibody is a mouse monoclonal antibody raised against TF. It has been selected for its ability to recognize TF in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig, Rat
Purification:	Protein A + Protein G affinity chromatography

## Target Details


Target:	Tissue factor (F3)
Alternative Name:	Tissue Factor ( <a href="#">F3 Products</a> )
Background:	CD142, FIII, F3, TFA, Thromboplastin, Coagulation Factor III
Pathways:	<a href="#">Positive Regulation of Endopeptidase Activity</a> , <a href="#">Smooth Muscle Cell Migration</a> , <a href="#">Platelet-derived growth Factor Receptor Signaling</a>

## Application Details

Application Notes:	Western blotting: 0.5-3 µg/mL Immunohistochemistry: 5-30 µg/mL Immunocytochemistry: 5-30 µg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months




#### Western Blotting

**Image 1.** Detection of TF in Rat Kidney lysate using Monoclonal Antibody to Tissue Factor (TF)



#### Immunohistochemistry

**Image 2.** Detection of TF in Human Cerebellum Tissue using Monoclonal Antibody to Tissue Factor (TF)



#### Immunohistochemistry

**Image 3.** Detection of TF in Human Pancreas Tissue using Monoclonal Antibody to Tissue Factor (TF)