antibodies - online.com







anti-ANG antibody (AA 24-148)





Overview

| Quantity: | 100 μL |
|----------------------|--|
| Target: | ANG |
| Binding Specificity: | AA 24-148 |
| Reactivity: | Cow |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This ANG antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details

| Purpose: | Monoclonal Antibody to Angiogenin (ANG) |
|---------------|---|
| Immunogen: | Recombinant Angiogenin (ANG) corresdonding to Ala24~His148 with N-terminal His Tag |
| Clone: | D4 |
| Isotype: | IgG |
| Specificity: | The antibody is a mouse monoclonal antibody raised against ANG. It has been selected for its ability to recognize ANG in immunohistochemical staining and western blotting. |
| Purification: | Protein A + Protein G affinity chromatography |


Target Details

Expiry Date:

| Target Details | |
|---------------------|---|
| Target: | ANG |
| Alternative Name: | Angiogenin (ANG Products) |
| Background: | RNASE5, Ribonuclease,RNase A Family 5 |
| Application Details | |
| Application Notes: | Western blotting: 0.5-5 μ g/mL Immunohistochemistry: 5-30 μ g/mL Immunocytochemistry: 5-30 μ g/mL Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without |

detectable loss of activity. Avoid repeated freeze-thaw cycles.

24 months



Western Blotting

Image 1. Detection of Recombinant ANG, Bovine using Monoclonal Antibody to Angiogenin (ANG)