

Datasheet for ABIN7427513 anti-CD163 antibody (AA 693-939)

2 Images



Overview

Quantity:	100 μL
Target:	CD163
Binding Specificity:	AA 693-939
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD163 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details


Purpose:	Monoclonal Antibody to Cluster Of Differentiation (CD163)
Immunogen:	Recombinant Cluster Of Differentiation (CD163) corresdonding to Pro693~Asp939 with N-terminal His Tag
Clone:	C8
Isotype:	IgG2a kappa
Specificity:	The antibody is a mouse monoclonal antibody raised against CD163. It has been selected for its ability to recognize CD163 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Human
Purification:	Protein A + Protein G affinity chromatography

Target Details

Expiry Date:


24 months

rarget Details	
Target:	CD163
Alternative Name:	Cluster Of Differentiation (CD163 Products)
Background:	M130, MM130, Scavenger receptor cysteine-rich type 1 protein M130, Hemoglobin scavenger receptor
Application Details	
Application Notes:	Western blotting: 0.5-5 μg/mL Immunohistochemistry: 5-30 μg/mL Immunocytochemistry: 5-30 μg/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.



Immunohistochemistry

Image 1. Detection of CD163 in Rat Stomach Tissue using Monoclonal Antibody to Cluster Of Differentiation (CD163)



Western Blotting

Image 2. Detection of Recombinant CD163, Rat using Monoclonal Antibody to Cluster Of Differentiation (CD163)