

Datasheet for ABIN7427810 **anti-Transferrin antibody**

1 Image



Go to Product page

| \sim | | | | |
|--------|----|-----|--------|---|
| ()\ | 10 | rVI | 161 | ٨ |
| \cup | ノロ | V | \Box | Λ |

| Quantity: | 100 μL |
|--------------|--|
| Target: | Transferrin (TF) |
| Reactivity: | Human |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This Transferrin antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

Product Details

Target:

| Purpose: | Monoclonal Antibody to Transferrin (TF) |
|----------------|---|
| Immunogen: | Native Transferrin (TF) |
| Clone: | 11# |
| Isotype: | IgG1 kappa |
| Specificity: | The antibody is a mouse monoclonal antibody raised against TF. It has been selected for its ability to recognize TF in immunohistochemical staining and western blotting. |
| Purification: | Protein A + Protein G affinity chromatography |
| Target Details | |

Transferrin (TF)

Target Details

| Alternative Name: | Transferrin (TF Products) |
|-------------------|---|
| Background: | TRF, Siderophilin, Serotransferrin, Beta-1 metal-binding globulin |
| UniProt: | P02787 |
| Pathways: | Transition Metal Ion Homeostasis |

Application Details

| Application Details | |
|---------------------|---|
| Application Notes: | Western blotting: 0.01-3 μ g/mL,Immunohistochemistry: 5-20 μ g/mL,Immunocytochemistry: 5-20 μ g/mL,Optimal working dilutions must be determined by end user. |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |

| Format: | Liquid |
|--------------------|---|
| Concentration: | 1 mg/mL |
| Buffer: | 0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles. |
| Expiry Date: | 24 months |



Western Blotting

Image 1. Detection of TF in Human Serum using Monoclonal Antibody to Transferrin (TF)