antibodies - online.com







anti-GIP antibody





Overview

Quantity:	100 μL
Target:	GIP
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This GIP antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Western Blotting (WB), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details


Target:

GIP

Purpose:	Monoclonal Antibody to Gastric Inhibitory Polypeptide (GIP)
Immunogen:	OVA Conjugated Gastric Inhibitory Polypeptide (GIP)
Clone:	A2
Isotype:	IgG2a lambda
Specificity:	The antibody is a mouse monoclonal antibody raised against GIP. It has been selected for its ability to recognize GIP in immunohistochemical staining and western blotting.
Purification:	Protein A + Protein G affinity chromatography
Target Details	

Target Details

9	
Alternative Name:	Gastric Inhibitory Polypeptide (GIP Products)
Background:	Incretin hormone, Glucose Dependent Insulinotropic Peptide
Pathways:	Positive Regulation of Peptide Hormone Secretion, Peptide Hormone Metabolism, Hormone Activity, Regulation of Lipid Metabolism by PPARalpha, Lipid Metabolism
Application Details	
Application Notes:	Immunocytochemistry in formalin fixed cells: 5-30 μ g/mL Immunohistochemistry in formalin fixed frozen section: 5-30 μ g/mL Immunohistochemistry in paraffin section: 5-30 μ g/mL Enzyme-linked Immunosorbent Assay: 0.05-2 μ g/mL Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months



Immunohistochemistry

Image 1. Detection of GIP in Human Pancreas Tissue using Monoclonal Antibody to Gastric Inhibitory Polypeptide (GIP)