antibodies -online.com





anti-Pepsinogen A antibody (AA 63-295)

3 Images



Overview


Quantity:	100 μL
Target:	Pepsinogen A
Binding Specificity:	AA 63-295
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Pepsinogen A antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC)

Product Details

Purpose:	Monoclonal Antibody to Pepsinogen A (PGA)
Immunogen:	Recombinant Pepsinogen A (PGA) corresdonding to Val63~Ser295 with N-terminal His Tag
Clone:	C4
Isotype:	IgG2b kappa
Specificity:	The antibody is a mouse monoclonal antibody raised against PGA. It has been selected for its ability to recognize PGA in immunohistochemical staining and western blotting.
Cross-Reactivity:	Pig
Purification:	Protein A + Protein G affinity chromatography

Target Details

•	
Target:	Pepsinogen A
Abstract:	Pepsinogen A Products
Background:	PGA5, PG-A, Pepsinogen 5 Group I, Pepsinogen-5, Pepsin A-5
Application Details	
Application Notes:	Western blotting: 0.5-2 μg/mL
	Immunohistochemistry: 5-20 μg/mL
	Immunocytochemistry: 5-20 μg/mL
	Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated
	thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious
	degradation and precipitation were observed. The loss rate is less than 5% within the expiration
	date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	0.01M PBS, pH 7.4, containing 0.05 % Proclin-300, 50 % glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be
	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without
	detectable loss of activity. Avoid repeated freeze-thaw cycles.
Expiry Date:	24 months




Western Blotting

Image 1. Detection of PGA in Porcine Stomach lysate using Monoclonal Antibody to Pepsinogen A (PGA)

Western Blotting

Image 2. Detection of Recombinant PGA, Human using Monoclonal Antibody to Pepsinogen A (PGA)



Immunohistochemistry

Image 3. Detection of PGA in Human Stomach Tissue using Monoclonal Antibody to Pepsinogen A (PGA)