# antibodies -online.com





## anti-PIIINP antibody



**Images** 



Go to Product page

| ( ) | 1 / | $\sim$   | KI / | 110 | Νę    |
|-----|-----|----------|------|-----|-------|
|     | 1// | $\vdash$ | I \/ | 1 ← | ٠// ٢ |
|     |     |          |      |     |       |


| Quantity:    | 100 μL   |
|--------------|--|
| Target:      | PIIINP   |
| Reactivity:  | Human  |
| Host:        | Mouse  |
| Clonality:   | Monoclonal   |
| Conjugate:   | This PIIINP antibody is un-conjugated  |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC) |

#### **Product Details**

| Purpose:          | Monoclonal Antibody to Procollagen III N-Terminal Propeptide (PIIINP)   |  |
|-------------------|---|--|
| Immunogen:        | Recombinant Procollagen III N-Terminal Propeptide (PIIINP) corresdonding to Two N-terminal Tags, His-tag and SUMO-tag   |  |
| Clone:            | E9  |  |
| Isotype:          | IgG2b kappa   |  |
| Specificity:      | The antibody is a mouse monoclonal antibody raised against PIIINP. It has been selected for its ability to recognize PIIINP in immunohistochemical staining and western blotting. |  |
| Cross-Reactivity: | Pig   |  |
| Purification:     | Protein A + Protein G affinity chromatography   |  |


## **Target Details**

| g                   |  |  |
|---------------------|--|--|
| Target:             | PIIINP   |  |
| Alternative Name:   | Procollagen III N-Terminal Propeptide (PIIINP Products)  |  |
| Background:         | P3NP, N-Propeptide Of Type III Procollagen, Procollagen III Amino Terminal Propeptide            |  |
| Application Details |  |  |
| Application Notes:  | Western blotting: 0.5-3 μg/mL  |  |
|                     | Immunohistochemistry: 5-30 μg/mL   |  |
|                     | Immunocytochemistry: 5-30 μg/mL  |  |
|                     | Optimal working dilutions must be determined by end user.  |  |
| Comment:            | The thermal stability is described by the loss rate. The loss rate was determined by accelerated |  |
|                     | thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious          |  |
|                     | degradation and precipitation were observed. The loss rate is less than 5% within the expiration |  |
|                     | date under appropriate storage condition.  |  |
| Restrictions:       | For Research Use only  |  |
| Handling            |  |  |
| Format:             | Liquid   |  |
| Buffer:             | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.                                      |  |
| Preservative:       | Sodium azide   |  |
| Precaution of Use:  | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which                    |  |
|                     | should be handled by trained staff only.   |  |
| Storage:            | 4 °C,-20 °C  |  |
| Storage Comment:    | Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without  |  |
|                     | detectable loss of activity. Avoid repeated freeze-thaw cycles.                                  |  |
| Expiry Date:        | 24 months  |  |
|                     |  |  |




### **Western Blotting**

**Image 1.** Detection of PIIINP in Rat Pancreas lysate using Monoclonal Antibody to Procollagen III N-Terminal Propeptide (PIIINP)



#### **Immunohistochemistry**

**Image 2.** Detection of PIIINP in Human Liver cancer Tissue using Monoclonal Antibody to Procollagen III N-Terminal Propeptide (PIIINP)



## **Immunohistochemistry**

**Image 3.** Detection of PIIINP in Human Colorectal cancer Tissue using Monoclonal Antibody to Procollagen III N-Terminal Propeptide (PIIINP)

Please check the product details page for more images. Overall 6 images are available for ABIN7428533.